NAMES.

A.

G. Κοκκορον

L. Fragaria.

I. Fravola.

G. Erdbeeren.

F. Fraises.

S. Miruedanos.

D. Aardbeeze

STRAWBERRIES.

PARTS USED.
Roots Leaves and Fruit.

PREPARATIONS.
None kept in the Shops.



PLACE.
They are cultivated in Gardens; some grow wild in many Places.

TIME.
Flower in May, Fruit ripe June and July.

DESCRIPTION.

The Roots are small, reddish, full of Fibres, that from Runners above Ground shoot Fibres downward, and Leaves upward, of a middling Green, Jagg'd, whiter underneath. The Stalks rise to support the Fruit above the Leaves, which from three upon one Foot-Stalk, abate to one near the Fruit. The Empalement 2 is a very light whitish Green set with white Hairs, divided into ten pointed Parts. The Flowers 3 have five white Leaves, in the Center of which is the young Fruit: at 4 'tis cut through to shew the Embrio of the Fruit, with the Seeds on the Outside, with every one a Yellow Pistil, with reddish Yellow Tops; the Stiles are white, with Summits cover'd with a Yellow Farina; the ripe Fruit has many light-colour'd whitish Seeds on their Outside, one of each Sort is over the Fruit that produced it. The Scarlet Strawberry 5, the Wood 6, which is a deeper Red, and the Hisboy 7, of which there are two Sorts, one Red or Black, the other White: 8 shews a Fruit cut Longitudinally, 9 another Transversly; the Inside is whitish.

VIRTUES.

The Leaves are used in Gargarisms, for Soreness of the Mouth &c. The Fruit is more used at Table than in Shops.

T. Sheldrake delin. R. Homerick sculp.